

Each drug is recommended for use in combination with other antiretrovirals (except any single-tablet regimen, including Atripla, Biktarvy, Complera, Genvoya, Juluca, Odefsey, Stribild, and Triumeq). For complete description of DHHS drug recommendations, go to [aidsinfo.nih.gov/guidelines](http://aidsinfo.nih.gov/guidelines). Consult with a physician about which drug therapy and dose is appropriate for you; see drug page or package insert for complete information. Drug chart information is current as of March 9, 2018. Look up your drug by adding its name after you type [positivelyaware.com/](http://positivelyaware.com/) into your browser (for example, [positivelyaware.com/triumeq](http://positivelyaware.com/triumeq)). For the complete POSITIVELY AWARE 2018 HIV Drug Guide, go to [positivelyaware.com](http://positivelyaware.com). For a list of Walgreens pharmacies that specialize in HIV/AIDS care and other resources, go to [HIV.walgreens.com](http://HIV.walgreens.com).

## SINGLE-TABLET REGIMENS (Multiple drug classes)



### Atripla RECOMMENDED INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

**efavirenz/emtricitabine/tenofovir DF (EFV/FTC/TDF)**  
One tablet (600 mg efavirenz/200 mg emtricitabine/300 mg tenofovir DF), once daily. Take on an empty stomach, preferably at bedtime.



### Complera RECOMMENDED INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

**rilpivirine/emtricitabine/tenofovir DF (RPV/FTC/TDF)**  
One tablet (25 mg rilpivirine/200 mg emtricitabine/300 mg tenofovir DF), once daily. Take with a meal.



### Juluca CAN BE A REASONABLE OPTION WHEN SWITCHING REGIMENS AND VIRALLY SUPPRESSED, THE USE OF NRTIS IS NOT DESIRABLE AND RESISTANCE TO EITHER DTG OR RPV IS NOT EXPECTED

**dolutegravir/rilpivirine (DTG/RPV)**  
One tablet (50 mg dolutegravir/25 mg rilpivirine) once daily. Take with a meal. For adults who are virologically suppressed on a current ART regimen for at least 6 months and who have no history of treatment failure or resistance mutations associated with rilpivirine or dolutegravir.



### Stribild RECOMMENDED INITIAL REGIMEN FOR MOST PEOPLE

**elvitegravir/cobicistat/emtricitabine/tenofovir DF (EVG/COBI/FTC/TDF)**  
One tablet (150 mg elvitegravir/150 mg cobicistat/200 mg emtricitabine/300 mg tenofovir DF), once daily. Take with food.



### darunavir/cobicistat/emtricitabine/tenofovir alafenamide (DRV/COBI/FTC/TAF) DHHS RECOMMENDATION NOT YET ESTABLISHED

One tablet (800 mg darunavir/50 mg cobicistat/200 mg emtricitabine/10 mg tenofovir alafenamide) once daily for patients without darunavir-related drug resistance. Take with food. (Dose cited from that used in EMERALD registrational study.)  
**NOT YET APPROVED AT PRESS TIME. PHOTO UNAVAILABLE.**



### Biktarvy RECOMMENDED INITIAL REGIMEN FOR MOST PEOPLE

**bictegravir/emtricitabine/tenofovir alafenamide (BIC/FTC/TAF)**  
One tablet (50 mg bictegravir/200 mg emtricitabine/25 mg tenofovir alafenamide) once daily. Take with or without food.



### Genvoya RECOMMENDED INITIAL REGIMEN FOR MOST PEOPLE

**elvitegravir/cobicistat/emtricitabine/tenofovir alafenamide (EVG/COBI/FTC/TAF)**  
One tablet (150 mg elvitegravir/150 mg cobicistat/200 mg emtricitabine/10 mg tenofovir alafenamide) once daily. Take with food.



### Odefsey RECOMMENDED INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

**rilpivirine/emtricitabine/tenofovir alafenamide (RPV/FTC/TAF)**  
One tablet (25 mg rilpivirine/200 mg emtricitabine/25 mg TAF) once daily. Take with a meal.



### Triumeq RECOMMENDED INITIAL REGIMEN FOR MOST PEOPLE

**dolutegravir/abacavir/lamivudine (DTG/ABC/3TC)**  
One tablet (50 mg dolutegravir/600 mg abacavir/300 mg lamivudine) once daily for people on HIV therapy for the first time. One additional 50 mg tablet of Tivicay 12 hours apart in people with viral resistance to INSTIs or when taken with certain other medications. Take with or without food.



### doravirine/tenofovir DF/lamivudine (DOR/TDF/3TC) DHHS RECOMMENDATION NOT YET ESTABLISHED

One tablet (100 mg/doravirine/300 mg lamivudine/300 mg tenofovir DF) once daily. Take with or without food. (Dose cited from that used in DRIVE-AHEAD registrational study.)  
**INVESTIGATIONAL DRUG AT PRESS TIME. PHOTO UNAVAILABLE.**

## INTEGRASE INHIBITORS



### Isentress HD RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

**and Isentress raltegravir (RAL)**  
Two 600 mg film-coated tablets once daily; twice daily Isentress available. See HIV Drug Guide for details. Take with or without food.



### Tivicay RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

**dolutegravir (DTG)**  
One 50 mg tablet once daily for people on HIV therapy for the first time or treatment-experienced without previous INSTI resistance. Twice-daily dosing for people who have viral resistance to INSTIs or when taken with certain other medications. Take with or without food.

## PROTEASE INHIBITORS (Boosted and unboosted)



### Evotaz RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

**atazanavir/cobicistat (ATV/COBI)**  
One tablet (300 mg atazanavir/150 mg cobicistat) once daily. Take with food.



### Prezcobix RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

**darunavir/cobicistat (DRV/COBI)**  
One tablet (800 mg darunavir/150 mg cobicistat) once daily, in patients with no darunavir drug resistance. Take with food.



### Prezista RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

**darunavir (DRV)**  
One 800 mg tablet with 100 mg Norvir or 150 mg Tybost, once daily for first-time therapy and treatment-experienced adults without Prezista-related resistance; or one 600 mg tablet plus 100 mg Norvir, twice daily for treatment-experienced people with Prezista-related resistance. Prezista must be taken with Norvir or Tybost. Take with food.



### Reyataz RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

**atazanavir sulfate (ATV)**  
One 300 mg capsule plus 100 mg Norvir or 150 mg Tybost, once daily; or two 200 mg capsules (without boosting), once daily for treatment-naïve adults (see the POSITIVELY AWARE HIV Drug Guide for details). Take with food. **GENERIC IS AVAILABLE.**

## PK ENHANCERS (Boosters)



### Tybost USED ONLY AS A BOOSTER FOR OTHER DRUGS; RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE AS WELL AS IN CERTAIN CLINICAL SITUATIONS

**cobicistat (COBI)**  
Used only for boosting—not an antiretroviral. 150 mg once daily with food taken at the same time with either Prezista 800 mg or Reyataz 300 mg. Take with food.



### Norvir USED ONLY AS A BOOSTER FOR OTHER DRUGS; RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS

**ritonavir (RTV)**  
Used only for boosting, also an antiretroviral. 100–200 mg, dosed once or twice daily with another PI. Must be taken with food. **GENERIC IS AVAILABLE.**

## NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS (“Nukes”)



### Descovy RECOMMENDED AS A COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

**emtricitabine/tenofovir alafenamide (FTC/TAF)**  
One tablet (200 mg emtricitabine/25 mg tenofovir alafenamide) once daily. Take with or without food.



### Emtriva RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

**emtricitabine (FTC)**  
One 200 mg capsule, once daily. Take with or without food.



### Epivir RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE WHEN USED IN COMBINATION WITH DOLUTEGRAVIR AND ABACAVIR (AS TRIUMEQ)

**lamivudine (3TC)**  
One 300 mg tablet, once daily; or one 150 mg tablet, twice daily. Take with or without food. **GENERIC IS AVAILABLE.**



### Epzicom RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE WHEN USED IN COMBINATION WITH DOLUTEGRAVIR (AS TRIUMEQ)

**abacavir/lamivudine (ABC/3TC)**  
One tablet (600 mg abacavir/300 mg lamivudine), once daily. Take with or without food. **GENERIC IS AVAILABLE.**



### Truvada RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

**emtricitabine/tenofovir DF (FTC/TDF)**  
One tablet (200 mg emtricitabine/300 mg tenofovir DF), once daily. Take with or without food. **APPROVED AS GENERIC; NOT YET COMMERCIALY AVAILABLE.**



### Viread RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE

**tenofovir disoproxil fumarate (TDF)**  
One 300 mg tablet, once daily. Take with or without food. **GENERIC IS AVAILABLE.**



### Ziagen RECOMMENDED AS COMPONENT OF INITIAL REGIMEN FOR MOST PEOPLE WHEN USED IN COMBINATION WITH DOLUTEGRAVIR AND LAMIVUDINE (AS TRIUMEQ)

**abacavir sulfate (ABC)**  
Two 300 mg tablets, once daily; or one 300 mg tablet, twice daily. Take with or without food. **GENERIC IS AVAILABLE.**

## NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS (“Non-nukes”)



### doravirine (DOR) DHHS RECOMMENDATION NOT YET ESTABLISHED

One tablet (100 mg of doravirine) once daily. Take with or without food. (Dose cited from that used in DRIVE-FORWARD registrational study.)  
**INVESTIGATIONAL DRUG AT PRESS TIME. PHOTO UNAVAILABLE.**



### Edurant RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS IN COMBINATION WITH DESCOVY OR TRUVADA (AS ODEFSEY OR COMPLERA)

**rilpivirine (RPV)**  
One 25 mg tablet, once daily. Take with a meal.



### Intelence FOR TREATMENT-EXPERIENCED PATIENTS WITH VIRAL STRAINS RESISTANT TO AN NRTI AND OTHER ANTIRETROVIRAL DRUGS ONLY

**etravirine (ETR)**  
One 200 mg tablet, twice daily. Take with food.



### Sustiva RECOMMENDED AS COMPONENT OF INITIAL REGIMEN IN CERTAIN CLINICAL SITUATIONS (AS A COMPONENT OF ATRIPLA, OR WITH DESCOVY OR TRUVADA)

**efavirenz (EFV)**  
One 600 mg tablet, once daily. Take on an empty stomach, preferably at bedtime. **GENERIC IS AVAILABLE.**

## ENTRY INHIBITOR



### Selzentry NOT RECOMMENDED AS COMPONENT OF AN INITIAL REGIMEN

**maraviroc (MVC)**  
150, 300, or 600 mg (available in 150 and 300 mg tablets), twice daily, depends on other medications used (see the POSITIVELY AWARE HIV Drug Guide for details). Take with or without food.

## CD4 POST-ATTACHMENT INHIBITOR Long-acting monoclonal antibody (mAb)



### Trozarzo DHHS RECOMMENDATION NOT YET ESTABLISHED

**ibalizumab-uiyk (IBA)**  
Administered once every two weeks via intravenous infusion. Treatment begins with an IV loading (starting) dose of 2,000 mg, followed by an 800 mg IV infusion maintenance dose given every two weeks thereafter.

## LEGACY DRUGS The following drugs are no longer or rarely prescribed:



**Aptivus**  
tipranavir (TPV)



**Combivir**  
lamivudine/zidovudine (3TC/AZT)



**Crixivan**  
indinavir (IDV)



**Fuzeon**  
enfuvirtide (T-20, or ENF)



**Invirase**  
saquinavir (SQV)



**Kaletra**  
lopinavir/ritonavir (LPV/r)



**Lexiva**  
fosamprenavir calcium (FPV)



**Rescriptor**  
delavirdine (DLV)



**Retrovir**  
zidovudine (AZT, or ZDV)



**Trizivir**  
abacavir/lamivudine/zidovudine (ABC/3TC/AZT)



**Videx EC**  
didanosine (ddl)



**Viracept**  
nelfinavir (NFV)



**Viramune XR**  
nevirapine (NVP)



**Zerit**  
stavudine (d4T)