

Each drug is recommended for use in combination with other antiretrovirals (except Atripla, Complera, Genvoya, Stribild, and Triumeq). For complete description of DHHS drug recommendations, go to aidsinfo.nih.gov/guidelines. Consult with a physician about which drug therapy and dose is appropriate for you; see drug page or package insert for complete information. Drug chart information is current as of February 5, 2016. Look up your drug by adding its name after you type positivelyaware.com/ into your browser (for example, positivelyaware.com/triumeq). For the complete POSITIVELY AWARE 2016 HIV Drug Guide, or for more information on drugs no longer recommended, go to positivelyaware.com. Special thanks to Christopher M. Nguyen, PharmD, for reviewing this chart. To view a list of Walgreens pharmacies that specialize in HIV/AIDS care and other resources, go to HIV.walgreens.com.

STR SINGLE-TABLET REGIMENS
INSTI INTEGRASE INHIBITORS
PI PROTEASE INHIBITORS
PKE PK ENHANCER
NRTI, NNRTI NUCLEOSIDE/NUCLEOTIDE REVERSE TRANSCRIPTASE INHIBITORS ("NUKES")
NNRTI NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS ("NON-NUKES")
LEGACY DRUGS (NO LONGER OR RARELY PRESCRIBED)

STR SINGLE-TABLET REGIMENS
INSTI INTEGRASE INHIBITORS
PI PROTEASE INHIBITORS
NRTI, NNRTI NUCLEOSIDE/NUCLEOTIDE REVERSE TRANSCRIPTASE INHIBITORS ("NUKES")
NNRTI NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS ("NON-NUKES")
E ENTRY INHIBITOR (CCR5 ANTAGONIST)
LEGACY DRUGS (NO LONGER OR RARELY PRESCRIBED)

Atripla efavirenz / emtricitabine / tenofovir, or EFV / FTC / TDF

- ✓ DHHS ALTERNATIVE

One tablet (600 mg efavirenz / 200 mg emtricitabine / 300 mg tenofovir), once daily. Take on an empty stomach, or with a light, low-fat snack.

Complera rilpivirine / emtricitabine / tenofovir DF, or RPV / FTC / TDF

- ✓ DHHS ALTERNATIVE ONLY IF HIV RNA < 100,000 C/ML AND CD4 > 200 CELLS/MM³

One tablet (25 mg rilpivirine / 200 mg emtricitabine / 300 mg tenofovir), once daily. Take with a meal.

Genvoya elvitegravir / cobicistat / emtricitabine / tenofovir alafenamide, or EVG / COBI / FTC / TAF

- ★ DHHS RECOMMENDED FOR FIRST-LINE USE

One tablet (150 mg elvitegravir / 150 mg cobicistat / 200 mg emtricitabine / 10 mg tenofovir alafenamide) once daily with food.

Stribild elvitegravir / cobicistat / emtricitabine / tenofovir DF, or EVG / COBI / FTC / TDF

- ★ DHHS RECOMMENDED FOR FIRST-LINE USE

One tablet (150 mg elvitegravir / 150 mg cobicistat / 200 mg emtricitabine / 300 mg tenofovir DF), once daily. Take with food.

Triumeq dolutegravir / abacavir / lamivudine, or DTG / ABC / 3TC

- ★ DHHS RECOMMENDED FOR FIRST-LINE USE

One tablet (50 mg dolutegravir / 600 mg abacavir / 300 mg lamivudine) once daily for people on HIV therapy for the first time. One additional 50 mg tablet of Tivicay 12 hours apart in people with viral resistance to Isentress or Vitekta (found in Genvoya and Stribild).

BRAND NAME NOT YET ESTABLISHED

 rilpivirine / emtricitabine / tenofovir alafenamide, or RPV / FTC / TAF

- DHHS RATING NOT YET ESTABLISHED

Dose to be established.

Isentress raltegravir, or RAL

- ★ DHHS RECOMMENDED FOR FIRST-LINE USE

One 400 mg tablet, twice daily. No food restrictions (take with or without food).

Tivicay dolutegravir, or DTG

- ★ DHHS RECOMMENDED FOR FIRST-LINE USE

One 50 mg tablet once daily for people on HIV therapy for the first time or treatment-experienced without previous INSTI resistance. Twice-daily dosing for people who have viral resistance to INSTIs and when taken with certain ARVs. No food restrictions (take with or without food).

Evotaz atazanavir / cobicistat

- ✓ DHHS ALTERNATIVE

One tablet (300 mg atazanavir/150 mg cobicistat) once daily with food. Use in treatment-experienced patients depends on protease inhibitor drug resistance substitutions.

Kaletra lopinavir / ritonavir, or LPV / r

- ➔ DHHS OTHER

Four tablets (200 mg lopinavir / 50 mg ritonavir), once daily if fewer than 3 lopinavir resistance mutations; or two 200 / 50 mg tablets, twice daily. Half-strength film-coated 100 mg lopinavir / 25 mg ritonavir tablet available. No food restrictions, but food may improve Norvir tolerability.

Norvir ritonavir, or RTV

- ★ USED ONLY AS A BOOSTER FOR OTHER DRUGS

Used mostly for boosting, 100–200 mg, dosed once or twice daily with another PI. Must be taken with food.

Prezcobix darunavir / cobicistat

- ✓ DHHS ALTERNATIVE

One tablet (800 mg darunavir/150 mg cobicistat) once daily with food, in patients with no darunavir drug resistance.

Prezista darunavir, or DRV

- ★ DHHS RECOMMENDED FOR FIRST-LINE USE

One 800 mg tablet with 100 mg Norvir or 150 mg Tybost, once daily for first-time therapy and treatment-experienced adults without Prezista-related resistance; or one 600 mg tablet plus 100 mg Norvir, twice daily for treatment-experienced people with Prezista-related resistance. Prezista must be taken with Norvir or Tybost. Take with food.

Reyataz atazanavir sulfate, or ATV

- ✓ DHHS ALTERNATIVE

One 300 mg capsule plus 100 mg Norvir or 150 mg Tybost, once daily; or two 200 mg capsules (without boosting), once daily for treatment-naïve adults (see the Drug Guide for details). Take with food.

Tybost cobicistat, or COBI

- ✓ USED ONLY AS A BOOSTER FOR OTHER DRUGS

Used only for boosting—not an antiretroviral. 150 mg once daily with food taken at the same time with either Prezista 800 mg or Reyataz 300 mg.

Emtriva emtricitabine, or FTC

- ★ DHHS RECOMMENDED FOR FIRST-LINE USE (A COMPONENT OF TRUVADA)

One 200 mg capsule, once daily. No food restrictions (take with or without food).

Epivir lamivudine, or 3TC

- ★ DHHS RECOMMENDED (CAN BE USED INTERCHANGEABLY WITH FTC)

One 300 mg tablet, once daily; or one 150 mg tablet, twice daily. No food restrictions (take with or without food). Generic is available.

Epzicom abacavir / lamivudine, or ABC / 3TC

- ✓ DHHS ALTERNATIVE (UNLESS WITH TIVICAY)

One tablet (600 mg abacavir / 300 mg lamivudine), once daily. No food restrictions (take with or without food).

Truvada emtricitabine / tenofovir DF, or FTC / TDF

- ★ DHHS RECOMMENDED FOR FIRST-LINE USE

One tablet (200 mg emtricitabine / 300 mg tenofovir), once daily. No food restrictions (take with or without food).

Viread tenofovir disoproxil fumarate, or TDF

- ★ DHHS RECOMMENDED (A COMPONENT OF TRUVADA)

One 300 mg tablet, once daily. No food restrictions (take with or without food).

Ziagen abacavir sulfate, or ABC

- ✓ DHHS ALTERNATIVE (A COMPONENT OF EPZICOM) UNLESS IN TRIUMEQ

Two 300 mg tablets, once daily; or one 300 mg tablet, twice daily. No food restrictions (take with or without food).

BRAND NAME NOT YET ESTABLISHED

 emtricitabine / tenofovir alafenamide, or FTC / TAF

- DHHS RATING NOT YET ESTABLISHED

Dose to be established.

Edurant rilpivirine hydrochloride, or RPV

- ✓ DHHS ALTERNATIVE WITH LIMITATIONS (A COMPONENT OF COMPLERA) AND ONLY WITH TRUVADA

One 25 mg tablet, once daily. Take with food.

Intelence etravirine, or ETR

- ▼ ONLY FOR TREATMENT-EXPERIENCED INDIVIDUALS

One 200 mg tablet, twice daily; or two 100 mg tablets, twice daily. Take with food.

Sustiva efavirenz, or EFV

- ✓ DHHS ALTERNATIVE WITH LIMITATIONS (A COMPONENT OF ATRIPLA)

One 600 mg tablet or three 200 mg capsules, once daily. Take on an empty stomach, or may be taken with a light, low-fat snack.

Selzentry maraviroc, or MVC

- ▼ USED ONLY IN CERTAIN SITUATIONS

150, 300, or 600 mg (available in 150 and 300 mg tablets), twice daily, depends on other medications used (see the POSITIVELY AWARE HIV Drug Guide for details). No food restrictions (take with or without food).

The following drugs are no longer or rarely prescribed, and appear only in the online version of the [POSITIVELY AWARE HIV Drug Guide](#):

 Aptivus tipranavir, or TPV	 Combivir lamivudine / zidovudine, or 3TC / AZT	 Crixivan indinavir, or IDV	 Fuzeon enfuvirtide, T-20, or ENF	 Invirase saquinavir, or SQV	 Lexiva fosamprenavir calcium, or FPV	 Rescriptor delavirdine, or DLV
 Retrovir zidovudine, AZT, or ZDV	 Trizivir abacavir / lamivudine / zidovudine, or ABC / 3TC / AZT	 Videx EC didanosine, or ddl	 Viracept nelfinavir, or NFV	 Vitekta elvitegravir, or EVG (Seldom used as a stand-alone drug.)	 Viramune XR nevirapine, or NVP	 Zerit stavudine, or d4T